

A SELF-REPORTED WORK SAMPLING
TO ASSESS THE EMERGENCY DEPARTMENT'S COSTS

P. CREMONESI, E. DI BELLA, M. MONTEFIORI, L. PERSICO

A self-reported work sampling to assess the Emergency Department's costs

P. Cremonesi, E. di Bella, **M. Montefiori***, L. Persico

Very Preliminary Version
Not to be quoted

Keywords: *emergency department, self-reporting work sampling, standard cost*

1. Introduction

The Emergency Department (ED) is one of the most demanding among all the hospital's departments both in terms of economic resources consumption and programming. The literature suggests they are responsible for a large share of the overall hospital's admissions [Williams (1996), Cremonesi et al. (2010), Sartini et al. (2007)] and examinations (such as x-rays) [Williams (1996)]. In the current scenario of reduced resources devoted to health care providers, hospitals are forced to closely scrutinize the information referring to their EDs with particular reference to their costs. EDs must, along with the other departments and the whole health care system, pursue the cost containment goal in such a way that the overall quality and the level of services provided would not decrease. Therefore it turns to be crucial to define indices and criteria to evaluate the emergency units both in terms of efficiency and quality. To this extent, the ED activity has to be monitored and optimized in order to provide the best outcome in terms of quality of care subject to a budget constraint. Optimization of patient flow and bottleneck elimination is a viable way at policy maker disposal to decrease operational cost and boost the quality of care [Tyrance et al. (1996)].

Strictly correlated with quality is the problem of overcrowding: delays in the ED may have particularly dramatic outcomes for patients. ED congestion is the cause of a two-sided problem: on the one hand congestion affects the quality of care, on the other hand it affects, because of inefficiency in the production process, the cost control. Tyrance et al. (1996) analyse the way non-urgent patients affect the high ED costs. Their paper reaches the conclusion that non-urgent ED accesses are not responsible of high cost in the US EDs. The same conclusion is reached by Williams et al. (1996) who estimate the average and the marginal cost both for urgent and non-urgent patients and show that the money saving that could be gained (transferring non-urgent patients to other "less expensive" structure) is negligible. Bamezai et al. (2005) reach a different conclusion and they conclude that the ED activity should be re-organized because of the high marginal cost for non-urgent ED visits.

With reference to the quality of the services provided when overcrowding is detectable, Hoot et al. (2006) study the ED overcrowding in terms of the national health service's quality, and Hoot et al. (2007) define four different criteria to measure and forecast the ED overcrowding. Kulstad et al. (2010) find a positive correlation between overcrowding and therapeutic errors using the Edwin score. Also Pines and Yealy (2009) consider ED overcrowding responsible for deficiencies in terms both of quality and effectiveness of the treatments provided. To conclude the survey it deserves to quote the study of Rossile et al. (2008) suggesting that the optimization of the clinical pathway of patients can't be defined disregarding the

*DIEM, University of Genoa, Italy montefiori@unige.it

Marcello Montefiori wish to thank the MIUR (within the FIRB project N. RBFRO8IKSB) for the funds received

elderly: an efficient programming of clinical pathway can't be done disregarding the peculiarities of people over 75.

This paper is intended to move in the direction of the analysis of ED cost composition and impact by a investigation based on microdata referring to the health related services provided by a ED belonging to an Italian primary Regional Hospital. The clinical data are matched up with the relevant accounting and economic information concerning the cost faced by the ED aiming at the goal of providing a new approach in order to identify the standard production costs and their variability between the different types of patients.

2. The database

In order to investigate Emergency Department cost structure, data referring to patient logistics have been collected along a whole week¹ and matched up with the relevant accounting and economic information concerning the cost incurred by the ED. By the electronic data processing center (W:OOD) of the ED of *Ospedali Galliera*, we have collected data referring to 1,045 patients. The information available with reference to each patient concerns: i. *Date and time of arrival* (it refers to triaging time); ii. *Medical attendant* (that is the identification code of the accepting medical staff); iii) *Triage entrance code*² (patient assignment to the classification code based on patient severity, it refers to the following severity scale: 0 = white, 1 = green, 2 = yellow; 3 = red; 4 = black i.e. deceased); iv. Patient's *personal information* (gender, date of birth, residence and nationality); v. *Means of transport*: arrival mode (by his own, ambulance, air ambulance, etc.); vi. *Admission cause* (morbidity, accident at work, traffic collision, violence, domestic accident, dog attack, scholastic injury, self-mutilation, biological accident, etc.); vii. *Arrival typology* (direct arrival, family doctor, specialist, emergency medical service, other hospital ward, etc.); viii. *Date and hour of visit* (that is the time of the first visit corresponding to patient assignment to a physician); ix. *Laboratory and non-laboratory prescription* (a separate dataset containing all the information about patient health treatment, as for instance, number of events, prescription code, date and hour of each event); x. *Event cost table* (it consists on a separate dataset containing, with reference to each prescription, all the relevant information about health treatment costs; xi. *Patient outcome* (ED healthcare process outcome: discharged, hospitalized, transferred to other medical structure, moved to OBI³ ward, moved to short hospitalization DB⁴ ward, abandonment, expelled, deceased in ED, dead on arrival, hospitalization refusing); xii. *Attending Physician* (identification code of the discharging doctor); xiii. *Prognosis* (number of prognosis days); xiv. *Discharging code* (patient re-assignment, when necessary, of a new triage code, i.e., at the end of the process it may be required to vary the triage code previously assigned); xv. *Date and hour of discharging* (it refers to the patient report closing time).

3. Methodology and data analysis

The main item in the ED balance sheet is the Medical staff cost whose imputation to each of the four Triage Codes is not automatically possible. A first approximation suggests to split the Medical staff cost proportionally to the time each doctor spends in visiting each patient, which in turn is characterized by a specific triage code. With this task in mind, we asked the Medical staff of the ED of E.O. Ospedali Galliera, a primary general hospital in Genova, to record the actual time that each ED Doctor dedicated to each

¹ From Thursday 9th December 2010, 8:00 pm to Thursday 16th December 2010, 8:00 pm.

² We refer to severity scale used in Italy in order to assess the overall severity of an ED admitted patient's illness based on the patient's unique clinical conditions, their interaction, and the resultant combined risk of morbidity and mortality.

³ Osservazione Breve Intensiva: hospital ward which admits patients with prognosis lower than 24 hours.

⁴ Degenza Breve: hospital ward which admits patients with prognosis greater than 24 hours but lower than 72 hours.

patient. Some basic statistics for the 1,045 patients visited during this week are given in Table 1. Consistently with the results shown in Cremonesi et al. (2010), red and yellow codes are older than white and green codes whereas the gender composition is quite balanced between males and females.

Table 1 - Descriptive statistics for the 1,045 patients visited by the ED Medical Personnel during the sample week.

Triage Code	Patients	Age (mean)	Gender (% male)
W	98	40	43%
G	706	45	51%
Y	219	64	48%
R	22	76	55%
Total	1,045	49	49%

At a deeper analysis level of the data collected, a significant difference in the visiting time emerged, because of a number of reasons. First, red codes require, generally, a longer visiting time, but at the same time there are some red codes (for example people affected by heart attacks) which do not require a long visiting time as the symptoms are easily identifiable and standard procedures are well defined in protocols. The twenty-one doctors cycled during the week and the number of patients they visited was variable according to the turns in which they worked during the week (workdays, week-end), hour of the day (day, night) and specialization (surgery, clinical doctors). It also happened that a few doctors had to visit and assist a few red or yellow codes whereas other doctors visited a lot of white and green codes. Therefore the variable "Medical Doctor" is relevant as each doctor has, for various reasons his own personal distribution of waiting times. Figure 1 gives an outlook through box-plots of all these waiting time distributions per each doctor. Moreover, in the rush of an ED where the life of people is in peril, it happened that some doctors forgot to note the visiting time of each patient so that only for 729 patients (70% of the total) this is available (see Table 2). Some patients have been visited by various doctors so that they have not been assigned to any of them and marked as NA.

Figure 1 - Box plots representing the visiting time distributions per doctor.




Table 2 - Visited patients and noted visiting times for each of the 21 doctors.

Doctor	Visited patients	Noted visiting times	%
D01	96	70	73%
D02	49	39	80%
D03	106	64	60%
D04	29	25	86%
D05	72	60	83%
D06	14	12	86%
D07	91	77	85%
D08	17	3	18%
D09	50	49	98%
D10	14	13	93%
D11	58	51	88%
D12	79	64	81%
D13	67	56	84%
D14	40	24	60%
D15	54	44	81%
D16	51	11	22%
D17	45	36	80%
D18	30	14	47%
D19	21	17	81%
D20	27	-	0%
D21	2	-	0%
NA	33	-	0%
Totals	1,045	729	70%

To face all the aforesaid points, and in particular to neutralize the "Medical Doctor" variable, a three steps analysis has been done:

1. all the visiting time distributions of the 21 physicians have been standardized in order to have distributions all having null mean and unitary variance and, *de facto*, neutralizing the difference in mean and dispersion among the different doctors;

2. On the basis of the standardized values, a linear model has been estimated to express the visiting time as a function of clinical and demographic variables (parentheses indicate the coefficient sign):

$$D_W = \beta_0 + \beta_1 D_G + \beta_2 D_Y + \beta_3 D_R + \beta_4 G + \beta_5 A + \beta_6 P_{nl} + \beta_7 P_l + \beta_8 OC + \beta_9 H + \beta_{10} FT \quad (1)$$

(−) (+) (+) (+) (+) (+) (+) (+) (−) (−) (−)

being:

- V_T : the standardized visiting time;
- D_G, D_Y, D_R : the triage color dummies (white triage color confounded with the intercept);
- G : Gender dummy (Male = 1; Female = 0);
- A : Patient's Age;
- P_{nl} : Number of Nolab Prescriptions;
- P_l : Number of Lab Prescriptions;
- OC : Overcrowding measured as number of patients that are contextually visited;
- H : Hospitalization Dummy (Hospitalized = 1; Not hospitalized = 0)
- FT : Fast Track Dummy (Fast Tracked patient = 1; Not Fast Tracked patient = 0)

The estimates of the parameters of model (1) and their significativity are given in Table 3.

3. On the basis of the econometric model derived, the 319 missing values have been estimated.

4. All the values have been de-standardized using a 5% trimmed mean and 5% trimmed variance in order to cut the 5% of extreme values (highest and lowest)

Table 3 - Regression coefficients for model (1).

Coefficients:	Estimate	Std. Error	t value	Pr(> t)	
<i>Intercept</i>	-0,83068	0,14590	-5,69300	0,00000	***
D_G	0,10037	0,11539	0,87000	0,38472	
D_Y	0,38601	0,14990	2,57500	0,01022	**
D_R	0,41014	0,24302	1,68800	0,09191	*
G	0,02925	0,05957	0,49100	0,62352	
A	0,00300	0,00157	1,91200	0,05622	*
P_{nl}	0,12431	0,01675	7,42000	0,00000	***
P_l	0,04377	0,00583	7,50500	0,00000	***
OC	-0,00476	0,00282	-1,68800	0,09183	*
H	-0,00348	0,09538	-0,03700	0,97087	
FT	-0,38333	0,12320	-3,11200	0,00194	***

Signif. codes: *** <0.01, ** <0.05, * <0.1

Residual standard error: 0.7893 on 715 degrees of freedom
 Multiple R-squared: 0.3864, Adjusted R-squared: 0.3778
 F-statistic: 45.02 on 10 and 715 DF, p-value: < 2.2e-16

The new waiting times for the triage codes are the ones given in Table 4. The Triage color weights have estimated computing the ratios between the color means and the white triage color mean (Table 4). For instance, white colors weight is 1 (4.57/4.57), green colors is 1.92 (8.78/4.57), yellow colors is 3.12 (14.26/4.57) and red colors is 3.41 (15/59/4.57). In other words it emerges that green codes have an average visiting time that is the double of the white codes while yellow and red codes have a visiting time that is more than three times the visiting time of white codes. Finally, yellow and red codes visiting times are one another much similar than white and green codes are.

Table 4: De-standardized means and Triage Color codes.

Triage Code	Patients	Minutes (mean)	Minutes (SD)	Weight
W	98	4.57	2.39	1.00
G	706	8.78	4.40	1.92
Y	219	14.26	5.63	3.12
R	22	15.59	7.11	3.41
Total	1,045	9.68	5.42	

If the mean visiting times are split into young people (people under 75 years old, with the dummy for Age equal to 0) and elderly (people over 75 with dummy for Age =1), the weights of color codes are given in Table 5. Although White codes have approximately the same weight independently from the age of the patient, for Green, Yellow and Red codes it comes out that the weight of elders is higher than the weight of younger even for the same severity level.

Table 5: De-standardized means and Triage Color codes for aged (AGE = 1) and young (AGE = 0) patients.

Triage Code	AGED	YOUNG
W	0.97	1.00
G	2.41	1.78
Y	3.25	2.93
R	3.56	2.67

4. Cost Analysis

The Emergency department faces different costs according to the structure we are observing. The table below provides the percentage composition of different cost type.

Table 6: Cost Composition

	Emergency Dep	Observation	Short Hospitalization
Fixed Costs Composition			
Medical Doctors	36,78%	12,82%	12,09%
Nurses	22,66%	61,38%	31,94%
Other Personnel	10,56%	0,00%	18,37%
Mortgages and other expenses	5,69%	3,66%	7,27%
Administrative Staff	1,26%		
Cleaning	0,35%	0,00%	4,37%
Fixed Costs Total	77,30%	77,86%	74,04%
Variable Costs Composition			
Surgical & Medical devices	2,68%	1,84%	2,18%
Drugs	1,78%	1,53%	1,84%
Kitchen & Laundry	0,35%	1,24%	4,41%
Health Services	0,36%		
Variable Costs Total	5,17%	4,61%	8,43%
Common Costs			
Common Costs	17,53%	17,53%	17,53%
TOTAL	100%	100%	100%

Analysing economic data, it emerges that the cost for physicians, nurses and other personnel counts for the 71,27%, 74,20%, 62,40% of the total costs incurred respectively by the ED, Observation and Short Hospitalization. Therefore information concerning the amount of time workers spend on particular activities is crucial in order to correctly assess the ED per-patient cost. To this extent during the aforementioned week of survey all physicians practicing in the ED were asked to note down the actual time dedicated to each patient. The interviewees were asked to precisely identify the actual time dedicated to each patient: preliminary visit, report reading and discharge procedures⁵.

By this self-reported work-sampling procedure we could add a new variable crucial to our analysis: xvi. Actual visiting time (consisting in the time physicians dedicated to each patient) and get the weights of table 4.

However, if at first we want to check for each patient cost and use this information as benchmark, it is possible to simply divide the cost taking into account the number of patients:

$$\frac{FC + VC + CC}{1045} = 135.19\text{€} = \begin{cases} 103.79 \text{ (fixed cost)} \\ 7.70 \text{ (variable cost)} \\ 23.70 \text{ (common cost)} \end{cases}$$

⁵ The ED activity is organized on three shifts (8:00 a.m. – 2:00 p.m. – 8:00 p.m.). At the end of each shifts, all the gathered reports were collected and verified in order to minimize the risk of missing values.

Table 7: Patient outcome

<i>Triage Code</i>	<i>Patients</i>	<i>ED</i>	<i>OBI</i>	<i>DB</i>
W	98	96	2	0
G	706	664	29	13
Y	219	162	40	17
R	22	19	0	3
Total	1,045	941	71	33

Using Table 7 **Errore. L'origine riferimento non è stata trovata.** it is possible to define the patient cost according to his outcome:

	Emergency Department	Observation	Short Hospitalization
Per patient cost	€ 116,39	€ 105,56	€ 735,02

But costs should be imputed according to the actual resource consumption that is generally strictly correlated with the patient severity. Cremonesi et al. (2010) arbitrarily weight the patients according to the triage code. In fact they observe that time spent to receive the first visit after the triage attribution decreases as the as long as the patient severity (i.e., the triage code) increases. On the other hand the time elapsing between the first visit and the exit time increases with the triage classification. To this extent they assume that a larger time period to exit implies a larger resource consumption by the patient in terms of: i) medical and non-medical staff; ii) clinical tests; iii) drugs; iv) equipment; v) other.

In this work the “Basic Scenario” of Cremonesi et al. (2010) is overcome through the definition of new weights estimated using the data collected in the sample week (cfr table 4). As it clearly emerges from Table 4 the white code patient turns to be a sort of benchmark for the weight associated to other colors. For instance a green code is assumed to have a cost of 1.92 times greater than a white. A yellow code costs 3.12 times the white. In other terms we may think at equivalent patients, where the “equivalence” refers to cost. The hospitalization of a red code patient is equivalent, in terms of resource absorption, to 3.41 white code patients. This new scenario suggests a new cost classification which takes into account the triage classification.

At this stage two hypothesis are possible:

$$1st\ HP \quad structure_{PatientCost_{TriageCode}} = \frac{FC + CC}{N. of Pat.} + \frac{VC}{Eq. Pat.} * Weight_{TriageCode} \quad (1)$$

$$2nd\ HP \quad structure_{PatientCost_{TriageCode}} = \frac{FC + CC + VC}{Eq. Pat.} * Weight_{TriageCode} \quad (2)$$

By the first hypothesis (1) the new equivalent patients are used to assess only the variable cost component, while the other cost component are computed without weighting patients. The assumption underneath relies on the fact that only variable costs should be affected by the intensity of the clinical assistance whereas fixed costs do not vary.

The second hypothesis (2) moves from the observation that we are searching, from an economic point of view, the actual resource consumption by different types of patients. To this extent all the cost incurred by the ED are shared using the weighted patients criterion.

To implement these hypothesis the computation of equivalent patient according to the outcome is required.

Triage Code	ED eq pat	OBI eq pat	DB eq pat
W	96	2	0
G	1274,88	55,68	24,96
Y	505,44	124,8	53,04
R	64,79	0	10,23
Total	1941,11	182,48	88,23

Table 8: Total costs per triage color, structure and hypothesis

	Hypothesis 1			Hypothesis 2		
	Emergency Department	Observation	Short Hospitalization	Emergency Department	Observation	Short Hospitalization
Cost per Equivalent Patient	€ 113,29	€ 102,59	€ 696,24	€ 56,42	€ 41,07	€ 274,92

We can observe that the cost for a white code might vary from a minimum of 41,07 € to a maximum of 696,24 € depending on the weight, the cost computation methodology and on the outcome of his clinical pathway. The 696,24 value refers to the cost for an equivalent patient, under hypothesis 1, which is hospitalized. However no white codes are short hospitalized in the week we are considering. To this extent is possible to assess that the minimum cost that the short hospitalization has to cope with is that related to the green code. The green code patient costs 717,56 under hypothesis 1 and € 527,85 under hypothesis 2.

We are persuaded that a standard cost definition is the necessary tool in the direction of a prospective reimbursement scheme implementation (based on tariffs) also with reference to the ED activity. As the economic literature has shown, a prospective payment (based on standard cost) would be the most effective incentive to induce efficiency in health care provision.

3. Results and discussion

In the present paper a self-reported work-sampling was implemented in order to define a weight function to be applied to calculate the actual patient cost with reference to ED activity. This issue represents one of the most relevant aspect of originality of this work. It emerges a great variability “between and within” the different types of patients depending also on the outcome, the patient severity and the health treatment structure. Table 4 presents some results concerning the per patient cost that emerges from the two hypotheses of cost classification presented in the previous section.

We believe that this kind of analysis well fits the federalizing process that Italy is experiencing. In fact the federal reform is driving our Country toward a decentralized provision and funding of local public services. The health care services are “fundamental” under the provisions of the law that in turn implies that a standard cost has to be defined for its funding. The standard cost (as it is defined by the law) relies on the concepts of appropriateness and efficiency in the production of the health care service, assuming a standard quality level as target. The identification and measurement of health care costs is therefore a crucial task propaedeutic to health services economic evaluation. Various guidelines with different amount of details have been set up for costing methods which, however, are defined in simplified frameworks and

using fictitious data. This study is a first attempt to proceed, using real data, in the direction of a precise definition of the costs inherent to the emergency department activity.

Acknowledgements

The authors would like to thank E.O. Ospedali Galliera for the precious cooperation in providing the data.

References

- Bamezai A., Melnick G., Nawathe A. (2005). The cost of an emergency department visit and its relationship to emergency department volume, *Annals of Emergency Medicine* 45(5):483-490
- Cao P., Toyabe S.I., Akazawa K. (2006). Development of a Practical Costing Method for Hospital, *Tohoku Journal of Experimental Medicine* 208:213-224
- Cremonesi P., di Bella E., Montefiori M. (2010). Cost analysis of emergency department, *Journal of Preventive Medicine and Hygiene*, vol.51 n.4, 2010;
- Cremonesi P., di Bella E., Montefiori M. (2011). Tempi d'attesa e sovraffollamento delle strutture di primo soccorso: un'analisi empirica. *Simeu Journal*, forthcoming
- Hoot N., Aronsky D. (2006). An Early System for Overcrowding in the Emergency Department, *AMIA 2006 Symposium Proceedings* p:339
- Hoot N., Zhou C., Jones I., Aronsky D. (2007). Measuring and Forecasting Emergency Department Crowding in Real Time, *Annals of Emergency Medicine* 49(6):747-755
- Horwitz L.I., Green J., Bradley E.H. (2010). US Emergency Department Performance on Wait Time and Length of Visit, *Annals of Emergency Medicine* 55(2):133-141
- Kulstad E.B., Sikka R., Sweis R.T., Kelley K.M., Rzechula K.H. (2010). ED overcrowding is associated with an increased frequency of medication errors, *American Journal of Emergency Medicine* 28:304-309
- McCarthy M.L., Aronsky D., Jones I.D., et al. (2008) The Emergency Department Occupancy Rate: A simple Measure of Emergency Department Crowding?, *Annals of Emergency Medicine* 51(1):15-24
- McCarthy M.L., Zeger S.L., Ding R., et al. (2009) Crowding Delays Treatment and Lengthens Emergency, *Annals of Emergency Medicine* 54(4):492-503
- Pines J.M., Yealy D.M. (2009). Advancing the Science of Emergency Department Crowding: Measurement and Solutions 54(4): 511-513
- Rossille D., Cuggia M., Arnault A., Bouget J., Le Beux P. (2008). Managing an emergency department by analysis HIS medical data: a focus on elderly patient clinical pathways, *Health Care Management Science* 11:139-146
- Sartini, M., Cremonesi, P., Tamagno, R., Cristina, M.L., Orlando, P and Simeu Group (2007). Quality in emergency departments: a study on 3,285,440 admissions. *Journal of Preventing Medicine and Hygiene*; 48: 17-23
- Tyrance, P.H., Himmelstein, D.U. and Woolhandler, S. (1996). *US Emergency Department Costs: No Emergency*, *American Journal of Public Health*; 86(11): 1527-1531
- Williams R.M. (1996). The costs of visits to emergency departments, *The New England Journal of Medicine* 334:642-646